arXiv:nucl-th/9811076v2 27 Mar 1999

Proton-Proton Scattering Lengths from Effective Field Theory

Xinwei Kong[1] and Finn Ravndal[1]

Department of Physics and Institute of Nuclear Theory,
University of Washington, Seattle, WA 98195, U.S.A

Abstract: Using a recently developed effective field theory for the interactions of nucleons at non-relativistic energies, we calculate the Coulomb corrections to proton-proton scattering. Including the dimension-eight derivative interaction in PDS regularization scheme, we obtain a modified Jackson-Blatt relation for the scattering lengths which is found to be phenomenologically satisfactory. The effective range is not modified by Coulomb effects to this order in the calculation.

PACS: 03.65.Nk; 13.75.Cs; 25.40.C

During the last year Kaplan, Savage and Wise have proposed an effective field theory for the interactions of nucleons at low energies[1]. These are characterized by scattering lengths which are much larger than the natural hadronic scale which in these systems is set by the pion mass. This made it initially difficult to construct consistent power counting rules which are necessary in any effective theory calculation to ensure that all contributions to a given order are included. By the introduction of a new regularization scheme called Power Divergence Subtraction (PDS) which is an extension of the more common MS scheme, these problems were solved. A very similar scheme was proposed at the same time by Gegelia[2]. Very recently, Mehen and Stewart[3] have made this off-shell scheme (OS) more well-defined and shown that it is in fact equivalent to the PDS scheme.

Within this framework several applications to problems involving the deuteron have now been made[4][5]. The first attempts to describe three-nucleon systems with two neutrons have also been initiated[6]. In this way the properties of the triton nucleus can be investigated. But also systems like ^3He with two protons are of obvious interest to consider in this new approach. For this to succeed, one must know how to include the repulsive Coulomb force between the two protons. As a first step in this direction we will here consider proton-proton scattering at very low energies.

The effective Lagrangian for non-relativistic protons with mass M in the spin-singlet channel is

$$\mathcal{L}_0 = p^\dagger \left(i\partial_0 + \frac{\boldsymbol{\nabla}^2}{2M} \right) p - \frac{C_0}{2} (p\sigma_2 p)(p\sigma_2 p)^\dagger \tag{1}$$

when we only include the lowest order contact interaction parameterized by the coupling constant C_0. It corresponds to the singular potential $C_0\delta(\mathbf{r})$ which will affect interactions

[1]On leave of absence from Institute of Physics, University of Oslo, N-0316 Oslo, Norway

only in the S-wave. In addition, we must also include the static Coulomb repulsion between the protons. It has the effective strength $\eta = \alpha M/2p$ where p is the CM momentum of protons, and thus also becomes strong at very low energies. This can be done using the well-established formalism based upon the exact solutions of the Schrödinger equation in the Coulomb potential[7]. Denoting these by $\psi(\mathbf{r})$, the probability to find the protons at the same point in the initial or final state is reduced and given by $|\psi(0)|^2 = C_\eta^2$ where

$$C_\eta^2 = \frac{2\pi\eta}{e^{2\pi\eta} - 1} \tag{2}$$

is the Sommerfeld factor[7]. In a partial wave expansion of the full scattering amplitude[8], the total phaseshift is $\sigma_\ell + \delta_\ell^C$ where σ_ℓ is the pure Coulombic shift and δ_ℓ^C is the shift due to the strong interactions. If we denote this latter one in the S-wave by δ_{pp}^C, it is related to the corresponding strong amplitude $T_{SC}(p)$ by the standard partial wave expression

$$p\,(\cot\delta_{pp}^C - i) = -\frac{4\pi}{M}\frac{e^{2i\sigma_0}}{T_{SC}(p)} \tag{3}$$

The superscript C is a reminder to the fact that this is not the strong phaseshift in the absence of Coulomb effects. Instead it is the strong phaseshift modified by the Coulomb potential. It is only the Coulomb repulsion between the protons in the initial and final states that have been removed at this stage.

As has been known for a long time, $\cot\delta_{pp}^C$ in (3) does not have a regular effective range expansion[9]. One finds instead

$$p\left[C_\eta^2\left(\cot\delta_{pp}^C - i\right) + 2\eta H(\eta)\right] = -\frac{1}{a_{pp}^C} + \frac{1}{2}r_0\,p^2 + \ldots \tag{4}$$

where a_{pp}^C and r_0 is respectively the S-wave Coulomb-modified scattering length and effective range. They arise after removing the part of the amplitude described by the complex function[10]

$$H(\eta) = \psi(i\eta) + \frac{1}{2i\eta} - \ln(i\eta) \tag{5}$$

which represents Coulomb effects at short distances. Its imaginary part cancels the unitary term $\sim i$ in (4). The real part defines the function $h(\eta) = \mathrm{Re}\psi(i\eta) - \ln\eta$ which is more suitable for phenomenological analysis[11].

By this approach one were able to extract the two scattering parameters a long time ago, $a_{pp}^C = -7.82\,\mathrm{fm}$ and the effective range $r_0 = 2.83\,\mathrm{fm}$ [11]. The scattering length is in magnitude much smaller than for pn and nn scattering in the spin singlet channel where one finds respectively the values $a_{pn} = -23.75\,\mathrm{fm}$ and $a_{nn} = -18.8\,\mathrm{fm}$. The difference between these two latter numbers is well within the range determined by the breaking of isospin invariance in strong interactions[12]. For proton-proton scattering on the other hand it became clear that not all short-distance effects of the Coulomb potential were taken

out by the H-function in (4). By investigating the detailed behavior of the Schrödinger equation for the coupled problem of strong and Coulomb forces at small separations Jackson and Blatt[14] succeeded in finding an expression for the scattering length a_{pp} due to only the strong interactions,

$$\frac{1}{a_{pp}} = \frac{1}{a_{pp}^C} + \alpha M \left[\ln \frac{1}{\alpha M r_0} - 0.33 \right] \tag{6}$$

With the above value for the effective range r_0 we then find $a_{pp} = -17.0\,\mathrm{fm}$ which is very close to the nn scattering length. The term -0.33 in the above formula is found to be only weakly dependent on the exact form of the strong potential[13].

From the effective Lagrangian (1) we can now calculate the proton-proton scattering amplitude. It will consist of the sum of all chains of proton bubbles separated by the vertex with value C_0. Including the Coulomb potential corresponds to having static photon exchanges between all proton lines in these Feynman diagrams. The Coulomb interactions in the initial and final states just give the Sommerfeld factor (2). In perturbation theory each bubble will be replaced by an infinite sum of bubbles with zero, one, two and so on Coulomb interactions as shown in Fig.1.



Figure 1: The Coulomb-dressed bubble is a sum of bubbles with static Coulomb exchanges.

Since the scattering amplitude represented by the diagrams in Fig.2 forms a geometric series, it thus takes the value

$$T_{SC}(p) = C_\eta^2 \frac{C_0\, e^{2i\sigma_0}}{1 - C_0\, J_0(p)} \tag{7}$$

Here $J_0(p)$ is the value of one bubble which equals the amplitude for a pair of protons with energy $E = p^2/M$ to move from zero separation to zero separation. This is given exactly by the the Coulomb propagator $G_C(E; \mathbf{r}' = 0, \mathbf{r} = 0)$. In momentum space it becomes

$$J_0(p) = M \int \frac{d^3k}{(2\pi)^3} \frac{2\pi\eta(k)}{e^{2\pi\eta(k)} - 1} \frac{1}{p^2 - k^2 + i\epsilon} \tag{8}$$

When this result for the scattering amplitude is now used in (3) and (4), we see that both the phaseshift σ_0 and the Sommerfeld factor $C_0^2(\eta)$ cancel out. We are thus only left with



Figure 2: The proton-proton scattering amplitude is a sum of chains of Coulomb-dressed bubbles.

the evaluation of (8) which we do using the PDS regularization scheme[1] in $d = 3 - \epsilon$ dimensions and with a renormalization mass μ. It is ultraviolet divergent which shows up as an $1/\epsilon$ pole in the integral. This will be cancelled by counterterms which renormalize the coupling C_0 in (7) to $C_0(\mu)$. As a result, the finite part of the dressed bubble (8) is found to be

$$J_0^{finite}(p) = \frac{\alpha M^2}{4\pi}\left[\ln\frac{\mu\sqrt{\pi}}{\alpha M} + 1 - \frac{3}{2}C_E - H(\eta)\right] - \frac{\mu M}{4\pi} \tag{9}$$

where $C_E = 0.5772\cdots$ is Euler's constant. The last term here is the contribution from the special PDS pole in $d = 2$ dimensions. We see now that also the function $H(\eta)$ cancels out in (4). There is no contribution to the effective range r_0 to this order in the effective theory. Defining the μ-dependent strong scattering length

$$\frac{1}{a_{pp}(\mu)} = \frac{4\pi}{MC_0(\mu)} + \mu \tag{10}$$

as for pn and nn interactions[1], we thus see from (4) and (7) that it can be expressed in terms of the measured scattering length a_{pp}^C as

$$\frac{1}{a_{pp}(\mu)} = \frac{1}{a_{pp}^C} + \alpha M\left[\ln\frac{\mu\sqrt{\pi}}{\alpha M} + 1 - \frac{3}{2}C_E\right] \tag{11}$$

This relation has the exact same form as the Jackson-Blatt relation (6) where now the renormalization length $1/\mu$ enters instead of the effective range. But $a_{pp}(\mu)$ is not a physical quantity in the sense that it can be measured directly and will thus in general depend on the renormalization point μ. Coulomb effects on length scales $> 1/\mu$ have been removed from it. One would expect the coupling constant $C_0(\mu)$ to be within a few percent of the corresponding coupling constants for pn and nn scattering. Our result is non-perturbative both in the strong coupling and in the fine structure constant which is seen to enter in the combination $\alpha \ln \alpha$. This is a consequence of the Coulomb force becoming strong when the energy goes to zero.

Since we are not including pions in the present effective theory, we must have the renormalization mass in the range $|1/a_{pp}^C| < \mu \leq m_\pi$. Depending on the exact value we see that the Coulomb correction can actually become of the same magnitude as the strong interaction. In fact, since the scattering length a_{pp}^C is negative, it can have a very big effect on the hadronic scattering length a_{pp}. This is seen from the right-hand side of (11) which becomes zero when $\mu \simeq m_\pi$ as shown in Fig.3. It is not a very physical result, corresponding to the hadronic scattering length a_{pp} being infinite.

The exact numerical properties of our result is obviously dependent on the regularization scheme we use. As a check, we have also calculated the Coulomb-dressed bubble (8) using instead a momentum cutoff Λ as a regulator. It will then appear in the logarithmic term of (11) instead of μ and the numerical constants inside the parenthesis will be different. In fact, the new result can be obtained from (11) by letting $\mu \approx 0.56\Lambda$. This corresponds to a blowup of the hadronic scattering length for $\Lambda \approx 240\,\mathrm{MeV}$.



Figure 3: Dependence of the inverse scattering length on the renormalization point. Dashed curve gives result with only C_0 interaction, while the solid curve also includes the C_2 interaction.

From a physical point of view such an unstability in the result is not acceptable. Instead we should like to see the dependence on the regulator mass to be as small as possible. The obvious way to cure this instability is to include higher order interactions in the effective Lagrangian (1). In the S-wave channel there is only one such interaction,

$$\mathcal{L}_2 = \frac{1}{4} C_2 (p\sigma_2 \overleftrightarrow{\nabla}^2 p)(p\sigma_2 p)^\dagger + h.c. \tag{12}$$

where the operator $\overleftrightarrow{\nabla} = 1/2(\overrightarrow{\nabla} - \overleftarrow{\nabla})$. Kaplan, Savage and Wise[1] showed that it can be included in a perturbative way and gives directly the effective range r_0 of the proton. The resulting Feynman diagrams can be obtained from the original diagrams by replacing one C_0 vertex in each diagram in Fig.2 by a C_2 vertex in every possible way. We then find the following correction to the scattering amplitude

$$\delta T_{SC}(p) = -\frac{C_2 e^{2i\sigma_0}}{[1 - C_0 J_0(p)]^2} \psi^*(0) \nabla^2 \psi(0) + \ldots \tag{13}$$

when we neglect higher orders in the fine structure constant. It depends crucially on the properties of the Coulomb wavefunction at the origin. While $|\psi(0)| = C_\eta$, the double derivative is divergent which is easily seen from the Schrödinger equation. A regulated expression can be obtained from the exact wavefunction which we transform to momentum space. The resulting integration is then done in the PDS scheme. In this way we find

$$\psi^*(0) \nabla^2 \psi(0)|_{reg} = C_\eta^2 \left(-p^2 + \alpha M \mu + \frac{1}{2} \alpha^2 M^2 \right) \tag{14}$$

after a rather elaborate calculation. The term proportional to μ is again from a PDS pole. Dropping the higher order α^2 term, we thus have the following correction to the inverse scattering amplitude

$$\delta\left(\frac{1}{T_{SC}(p)}\right) = \frac{e^{-2i\sigma_0}C_2(\mu)}{C_\eta^2 C_0^2(\mu)}\left(p^2 - \alpha M\mu\right) \tag{15}$$

Comparing with the effective range expansion (4) we then obtain for the effective range of the proton simply $r_0 = 8\pi C_2/MC_0^2$. It is therefore not modified by the Coulomb interactions. This is consistent with experiments which show that the effective range for pp is the same as for pn and nn to within a few percent[12].

With the coupling constant C_2 determined in this way, we can now calculate its contribution to the scattering length by taking the limit of zero momentum in (15). The PDS pole is seen to give an extra term in the previous result (11) which now is changed into

$$\frac{1}{a_{pp}(\mu)} = \frac{1}{a_{pp}^C} + \alpha M\left[\ln\frac{\mu\sqrt{\pi}}{\alpha M} + 1 - \frac{3}{2}C_E - \frac{1}{2}\mu r_0\right] \tag{16}$$

Although this new term has been calculated in perturbation theory, it comes into the result with the same general magnitude as the first Coulomb correction and actually stabilizes the result when considered as a function of the renormalization point μ. This is shown in Fig.3 where the corresponding curve varies little in the whole interval $80\,\mathrm{MeV} < \mu < 200\,\mathrm{Mev}$. The blowup of the scattering length for μ approaching the pion mass has now been turned into a rather smooth behaviour in the same region. In fact, we find $-1/a(\mu = m_\pi) = 6.6\,\mathrm{MeV}$. For comparison, it should be noted that $-1/a_{pn} = 8.3\,\mathrm{MeV}$ and $-1/a_{nn} = 10.5\,\mathrm{MeV}$ while (6) gives $-1/a_{pp} = 11.6\,\mathrm{MeV}$. These differences in scattering lengths correspond to much smaller differences in the corresponding coupling constants C_0. The strong scattering length in our approach is scheme dependent and it will not enter explicittly in any physical results.

We would like to thank Martin Savage and all other members of the EFT group for much help and many useful discussions. In addition, we want to thank the Department of Physics and the INT for generous support and hospitality. Xinwei Kong is supported by the Research Council of Norway.

References

[1] D.B. Kaplan, M.J. Savage and M.B. Wise, *Phys. Lett.* **B424**, 390 (1998); *nucl-th/9802075.*

[2] J. Gegelia, *nucl-th/9802038*; *Phys. Lett.* **B429**, 227 (1998).

[3] T. Mehen and I.W. Stewart, *nucl-th/9809071, 9809095*

[4] D.B. Kaplan, M.J. Savage and M.B. Wise, *nucl-th/9804032*.

[5] J.-W. Chen, H.W. Griesshammer, M.J. Savage and R.P. Springer, *nucl-th/9806080, 9809023*.

[6] P.F. Bedaque and U. van Kolck, *Phys. Lett.* **B428**, 221 (1998); P. F. Bedaque, H.W. Hammer and U. van Kolck, *nucl-th/9809025*.

[7] A. Sommerfeld, *Atombau und Spektrallinien*, Vol. II, Vieweg, Braunschweig, 1939; L.D. Landau and E.M. Lifschitz, *Quantum Mechanics*, Pergamon Press, London, 1958.

[8] M.L. Goldberger and K.M. Watson, *Collision Theory*, John Wiley and Sons, New York, 1964; D.R. Harrington, *Phys. Rev.* **139**, B691 (1965).

[9] H.A. Bethe, *Phys. Rev.* **76**, 38 (1949).

[10] L.P. Kok, J.W. de Maag, H.H. Bouwer and H. van Haeringen, *Phys. Rev.* **C26**, 2381 (1982).

[11] G.E. Brown and A.D. Jackson, *The Nucleon-Nucleon Interaction*, North-Holland Publishing Company, Amsterdam, 1976.

[12] E.M. Henley in *Isospin in Nucelar Physics*, ed. D.H. Wilkinson, North-Holland Publishing Company, Amsterdam, 1969; G.A. Miller, B.M.K. Nefkens and I. Slaus, *Phys. Rep.* **194**, 1 (1990).

[13] L. Heller, P. Signell and N.R. Yoder, *Phys. Rev. Lett.* **13**, 577 (1964).

[14] J.D. Jackson and J.M. Blatt, *Rev. Mod. Phys.* **22**, 77 (1950).